

Very Differentiable and Generic Frechet Differentiable Convex Functions on Banach Spaces

Xi Yin Zheng

metadata, citation and similar papers at core.ac.uk

Submitted by William A. Kirk

Received December 10, 1997

Using the extension of convex functions on a Banach space X to the bidual space X^{**} , we introduce and study a kind of differentiability of convex functions. Moreover, we discuss Asplund spaces. © 1999 Academic Press

1. INTRODUCTION

Throughout this paper, X always denotes a Banach space. X^* , X^{**} , and X^{***} stand for the dual space, bidual space, and tridual space of X , respectively. We always regard X as a subspace of X^{**} . For $x \in X$, $B_X(x, r)$, and $B_{X^{**}}(x, r)$ stand for the ball with center x and radius r in X and X^{**} , respectively. For a subset A of X , we denote the weak* closure of A in X^{**} by $w^*\text{-cl}_{X^{**}}(A)$. w and w^* always stand for the weak topology and the weak* topology, respectively.

Let D be an open convex subset of X , $f: D \rightarrow \mathbb{R}$ a continuous convex function. For $x \in D$, $\partial f(x)$ denotes the subdifferential of f at x , that is,

$$\partial f(x) = \{x^* \in X^*; \langle x^*, y - x \rangle \leq f(y) - f(x), y \in D\}.$$

It is well known that ∂f is a $\|\cdot\|$ - w^* upper semicontinuous set-valued mapping and $\partial f(x)$ is a nonempty w^* compact convex subset of X^* for all $x \in D$. We say that f is Gateaux differentiable at $x \in D$ if the limit

$$\lim_{t \rightarrow 0} \frac{f(x + th) - f(x)}{t}$$



exists for all $h \in X$. This is equivalent to $\partial f(x)$ being a singleton. We say that f is Frechet differentiable at $x \in D$ if the limit

$$\lim_{t \rightarrow 0} \frac{f(x + th) - f(x)}{t}$$

exists uniformly with respect to h on the unit ball of X . It is known that f is Frechet differentiable at $x \in D$ iff $\partial f(x)$ is a singleton and ∂f is $\|\cdot\|$ -upper semicontinuous at x .

We say that X is an Asplund space if each continuous convex function defined on an open convex subset D of X is Frechet differentiable at each point in a dense G_δ subset of D . It is clear that X is an Asplund space iff for any open convex subset D of X and any continuous convex function f defined on D , the set $E = \{x \in D; \partial f(x) \text{ is a singleton and } \partial f \text{ is } \|\cdot\| \text{-upper semicontinuous at } x\}$ is a residual subset of D (i.e., E includes a dense G_δ subset of D). Asplund spaces are a very important subject in Banach spaces theory. A series of profound results on Asplund spaces were obtained (see [1, 6–12] and references therein).

In Section 2, we present some results about an extension of a continuous convex function defined on an open subset of a Banach space X to X^{**} . In Section 3, using the extension of convex functions, we introduce and study very differentiability of convex functions, which is a generalization of differentiability (or smoothness) of norms and, in general, is not any kind of β -differentiability. In Section 4, using the result that a continuous convex function f defined on an open convex subset D of a Banach space is Frechet differentiable at each point in a dense G_δ subset of D iff $D_f = \{x \in D; \partial f(x) \text{ is a weak compact set and } \partial f \text{ is } \|\cdot\| \text{-upper semicontinuous at } x\}$ is a residual subset of D , we give a characterization of Asplund spaces.

2. PRELIMINARIES

Let D be an open convex subset of X , f a continuous convex function defined on D . Pick a $x_0 \in D$ and $x_0^* \in \partial f(x_0)$; then $f(x) \geq f(x_0) + \langle x_0^*, x - x_0 \rangle$ for all $x \in D$. This implies that for each $x^{**} \in w^*\text{-cl}_{X^{**}}(D)$

$$\liminf_{x_D \xrightarrow{w^*} x^{**}} f(x) \geq f(x_0) + \langle x_0^*, x^{**} - x_0 \rangle > -\infty.$$

Hence we can define $\hat{f}: X^{**} \rightarrow R \cup \{+\infty\}$ such that

$$\hat{f}(x^{**}) = \begin{cases} \liminf_{x_D \xrightarrow{w^*} x^{**}} f(x), & x^{**} \in w^*\text{-cl}_{X^{**}}(D), \\ +\infty, & \text{otherwise.} \end{cases}$$

This means that the following (I) and (II) hold:

(I) If a net $\{x_\alpha\}$ in D converges to some $x^{**} \in w^*\text{-cl}_{X^{**}}(D)$ with respect to the weak* topology of X^{**} ,

$$\liminf_{\alpha} f(x_\alpha) \geq \hat{f}(x^{**})$$

(II) For each $x^{**} \in w^*\text{-cl}_{X^{**}}(D)$ there is a net $\{x_\alpha\}$ in D such that $x_\alpha \xrightarrow{w^*} x^{**}$ and

$$\hat{f}(x^{**}) = \lim_{\alpha} f(x_\alpha).$$

PROPOSITION 2.1. *Let f be a continuous convex function defined on an open convex subset D of X . Then $\text{epi}(\hat{f}) = w^*\text{-cl}_{X^{**} \times R}(\text{epi}(f))$. Hence \hat{f} is a proper weak* lower semicontinuous convex function on X^{**} .*

Proof. For any $(x^{**}, r) \in \text{epi}(\hat{f})$, $x^{**} \in w^*\text{-cl}_{X^{**}}(D)$ and there is $t \geq 0$ such that $r = f(x^{**}) + t$. Pick a net $\{x_\alpha\}$ in D such that $x_\alpha \xrightarrow{w^*} x^{**}$ and $f(x_\alpha) \rightarrow \hat{f}(x^{**})$. It follows that $(x_\alpha, f(x_\alpha) + t) \xrightarrow{w^*} (x^{**}, r)$ in $X^{**} \times R$. Hence $(x^{**}, r) \in w^*\text{-cl}_{X^{**} \times R}(\text{epi}(f))$. On the other hand, for any $(x^{**}, r) \in w^*\text{-cl}_{X^{**} \times R}(\text{epi}(f))$, there is a net $\{x_\alpha\}$ in D and a net $\{t_\alpha\}$ in R^+ such that $(x_\alpha, f(x_\alpha) + t_\alpha) \xrightarrow{w^*} (x^{**}, r)$. Hence

$$r = \lim_{\alpha} (f(x_\alpha) + t_\alpha) \geq \liminf_{\alpha} f(x_\alpha) \geq \hat{f}(x^{**}).$$

This implies $(x^{**}, r) \in \text{epi}(\hat{f})$. ■

PROPOSITION 2.2. *For each continuous convex function f defined on an open convex subset D of X , \hat{f} is an extension of f and $D \subset \text{int}_{X^{**}}(\text{dom}(\hat{f}))$, where $\text{dom}(\hat{f}) = \{x^{**} \in X^{**}; \hat{f}(x^{**}) \in R\}$.*

Proof. For each $x \in D$, by (I), $\hat{f}(x) \leq f(x)$. By (II) there is a net $\{x_\alpha\}$ in D such that $x_\alpha \xrightarrow{w^*} x$ (i.e., $\{x_\alpha\}$ converges to x with respect to the weak* topology of X^{**}) and $\hat{f}(x) = \lim_{\alpha} f(x_\alpha)$. Since a continuous convex function is lower semicontinuous with respect to the weak topology, $f(x) \leq \lim_{\alpha} f(x_\alpha) = \hat{f}(x)$. Hence $f(x) = \hat{f}(x)$ for all $x \in D$. By the conti-

nuity of f , for each $x \in D$ there are $\delta > 0$ and $M \in R$ such that $B_X(x, \delta) \subset D$ and $f(x) \leq M$ for all $x \in B_X(x, \delta)$. By the Goldstine theorem, $B_X(x, \delta)$ is weak* dense in $B_{X^{**}}(x, \delta)$. Hence, for each $x^{**} \in B_{X^{**}}(x, \delta)$, there is a net $\{z_\alpha\}$ in $B_X(x, \delta)$ such that $z_\alpha \xrightarrow{w^*} x^{**}$. It follows from (I) that $\hat{f}(x^{**}) \leq M$. Hence $x^{**} \in \text{dom}(\hat{f})$. This shows that $D \subset \text{int}_{X^{**}}(\text{dom}(\hat{f}))$. ■

LEMMA 2.1. *Let K be a nonempty bounded subset of X^* with $0 \in K$, and let $p_K(x) = \sup\{\langle x^*, x \rangle; x^* \in K\}$ for all $x \in X$ and $q_K(x^{**}) = \sup\{\langle x^{**}, x^* \rangle; x^* \in w^*\text{-cl}(\text{co}(K))\}$ for all $x^{**} \in X^{**}$. Then $A = \{x^{**} \in X^{**}; q_K(x^{**}) \leq 1\}$ is the weak* closure of $B = \{x \in X; p_K(x) \leq 1\}$.*

Proof. It is easy to verify that $p_K(x) = \sup\{\langle x^*, x \rangle; x^* \in w^*\text{-cl}(\text{co}(K))\}$ for all $x \in X$. Hence, $q_K|_X = p_K$. Notice that q_K is a positively homogeneous, subadditive, and weak* lower semicontinuous function on X^{**} . We have $w^*\text{-cl}_{X^{**}}(B) \subset A$. Suppose that there is $x_0^{**} \in A$ such that $x_0^{**} \notin w^*\text{-cl}_{X^{**}}(B)$. By the separation theorem, there is $x_0^* \in X^*$ and $r \in R$ such that

$$\langle x_0^{**}, x_0^* \rangle > r > \sup\{\langle x_0^*, x \rangle; x \in B\} \geq 0.$$

It follows that $\sup\{\langle \frac{1}{r}x_0^*, x \rangle; x \in B\} \leq 1$. By the separation theorem, it is easy to verify $\frac{1}{r}x_0^* \in w^*\text{-cl}(\text{co}(K))$. Hence $q_K(x_0^{**}) \geq \langle x_0^{**}, \frac{1}{r}x_0^* \rangle > 1$. This contradicts $x_0^{**} \in A$. ■

EXAMPLE 2.1. Let K be a nonempty bounded subset of X^* , and let $f(x) = \sup\{\langle x^*, x \rangle; x^* \in K\}$ for all $x \in X$. Then

$$\hat{f}(x^{**}) = \sup\{\langle x^{**}, x^* \rangle; x^* \in w^*\text{-cl}(\text{co}(K))\}, \quad x^{**} \in X^{**}.$$

In particular, when $f(x) = \|x\|$ for all $x \in X$, $\hat{f}(x^{**}) = \|x^{**}\|$ for all $x^{**} \in X^{**}$.

Proof. Let $g(x^{**}) = \sup\{\langle x^{**}, x^* \rangle; x^* \in w^*\text{-cl}(\text{co}(K))\}$ for all $x^{**} \in X^{**}$; then $g|_X = f$. By the weak* lower semicontinuity of g , $w^*\text{-cl}_{X^{**}}(X) = X^{**}$ and (II), $g(x^{**}) \leq \hat{f}(x^{**})$ for all $x^{**} \in X^{**}$. Suppose that there is $x_0^{**} \in X^{**}$ such that $g(x_0^{**}) < \hat{f}(x_0^{**})$. Pick a $x_0^* \in K$ and $r > 0$ such that

$$g(x_0^{**}) - \langle x_0^{**}, x_0^* \rangle < r < \hat{f}(x_0^{**}) - \langle x_0^{**}, x_0^* \rangle.$$

Let $A = \{x^{**} \in X^{**}; g(x^{**}) - \langle x^{**}, x_0^* \rangle \leq r\}$ and $B = \{x \in X; f(x) - \langle x_0^*, x \rangle \leq r\}$. By Lemma 2.1, $A = w^*\text{-cl}_{X^{**}}(B)$. Hence $x_0^{**} \in w^*\text{-cl}_{X^{**}}(B)$, and so there is a net $\{x_\alpha\}$ in B such that $x_\alpha \xrightarrow{w^*} x_0^{**}$. It follows that

$$\hat{f}(x_0^{**}) \leq \liminf_{\alpha} f(x_\alpha) \leq \langle x_0^{**}, x_0^* \rangle + r.$$

Hence $\hat{f}(x_0^{**}) - \langle x_0^{**}, x_0^* \rangle \leq r$, a contradiction. ■

3. VERY DIFFERENTIABLE CONVEX FUNCTIONS

It is an aspect of Banach spaces theory to study very smoothness of a space (see [11] and references therein). A point x in X with $\|x\| = 1$ is said to be a very differentiable point of X if there is a unique x^{***} in X^{***} such that $\|x^{***}\| = 1$ and $\langle x^{***}, x \rangle = 1$. This is equivalent to the norm of X^{**} being Gateaux differentiable at x . This and Example 2.1 incline us to introduce the very differentiability of convex functions.

DEFINITION 3.1. Let f be a continuous convex function defined on an open convex subset D of X . We say that f is very differentiable at $x \in D$ if \hat{f} , as a function on X^{**} , is Gateaux differentiable at x .

THEOREM 3.1. Let f be a continuous convex function defined on an open convex subset D of X , and $x_0 \in D$. Then for each $x^{***} \in \partial \hat{f}(x_0)$ there is a net $\{x_\alpha\}$ in D and a net $\{x_\alpha^*\}$ in X^* such that $x_\alpha \rightarrow x_0$, $x_\alpha^* \xrightarrow{w^*} x^{***}$ and for each index α , $x_\alpha^* \in \partial f(x_\alpha)$.

Proof. Let $C(f) = \{(x^*, r) \in X^* \times R; \langle x^*, x \rangle + r t \leq 1, (x, t) \in \text{epi}(f) - (x_0, f(x_0) + 1)\}$. We claim that for any $x^{***} \in \partial \hat{f}(x_0)$, $(x^{***}, -1) \in w^*\text{-cl}_{X^{***} \times R}(C(f))$. Indeed, suppose that there is $x_0^{***} \in \partial \hat{f}(x_0)$ such that $(x_0^{***}, -1) \notin w^*\text{-cl}_{X^{***} \times R}(C(f))$. By the separation theorem, there is $(x_0^{**}, r_0) \in X^{**} \times R$ and $t_0 \in R$ such that

$$\langle x_0^{***}, x_0^{**} \rangle - r_0 > t_0 > \sup\{\langle x_0^{**}, x^* \rangle + r_0 r; (x^*, r) \in C(f)\}.$$

Since $(0, 0) \in C(f)$, $t_0 > 0$. Hence

$$\left\langle x_0^{***}, \frac{1}{t_0} x_0^{**} \right\rangle - \frac{r_0}{t_0} > 1 \quad (1)$$

and

$$\sup\left\{\left\langle \frac{1}{t_0} x_0^{**}, x^* \right\rangle + \frac{r_0 r}{t_0}; (x^*, r) \in C(f)\right\} < 1.$$

This and the separation theorem imply

$$\left(\frac{1}{t_0} x_0^{**}, \frac{r_0}{t_0}\right) \in w^*\text{-cl}_{X^{**} \times R}(\text{epi}(f) - (x_0, f(x_0) + 1)).$$

By Proposition 2.1, $((1/t_0)x_0^{**}, (r_0/t_0)) \in \text{epi}(\hat{f}) - (x_0, f(x_0) + 1)$. Hence there is $s_0 \geq 0$ such that

$$\frac{r_0}{t_0} = \hat{f}\left(\frac{1}{t_0}x_0^{**} + x_0\right) + s_0 - f(x_0) - 1.$$

This and (1) imply

$$\begin{aligned} \left\langle x_0^{***}, \frac{1}{t_0}x_0^{**} \right\rangle &> \hat{f}\left(\frac{1}{t_0}x_0^{**} + x_0\right) + s_0 - f(x_0) \\ &\geq \hat{f}\left(\frac{1}{t_0}x_0^{**} + x_0\right) - \hat{f}(x_0). \end{aligned}$$

This contradicts $x_0^{***} \in \partial\hat{f}(x_0)$. Hence for any $x^{***} \in \partial\hat{f}(x_0)$, there is a net $\{(x_\alpha^*, r_\alpha)\}$ in $C(f)$ such that $(x_\alpha^*, r_\alpha) \xrightarrow{w^*} (x^{***}, -1)$ in $X^{***} \times R$, that is, $x_\alpha^* \xrightarrow{w^*} x^{***}$ in X^{***} and $r_\alpha \rightarrow -1$. By $(x_\alpha^*, r_\alpha) \in C(f)$, for each $x \in D$,

$$\langle x_\alpha^*, x - x_0 \rangle + r_\alpha(f(x) - f(x_0) - 1) \leq 1,$$

that is,

$$\langle x_\alpha^*, x - x_0 \rangle \leq f(x) - f(x_0) + (1 + r_\alpha)(f(x_0) - f(x) + 1). \quad (2)$$

By the continuity of f , there is $\delta > 0$ such that $B_X(x_0, \delta) \subset D$ and $|f(x) - f(x_0)| < 1$ for all $x \in B_X(x_0, \delta)$. Let $\varepsilon_\alpha = 2|1 + r_\alpha|$; then $\varepsilon_\alpha \rightarrow 0$. Setting $g = f|_{B_X(x_0, \delta)}$, by (2), $\langle x_\alpha^*, x - x_0 \rangle \leq g(x) - g(x_0) + \varepsilon_\alpha$ for each $x \in B_X(x_0, \delta)$. Hence $x_\alpha^* \in \partial_{\varepsilon_\alpha} g(x_0)$. By the Brondsted-Rockafellar theorem (see [10, p. 51, Theorem 3.18], there are $x_\alpha \in B_X(x_0, \delta)$ and $y_\alpha^* \in \partial g(x_\alpha) = \partial f(x_\alpha)$ such that

$$\|x_\alpha - x_0\| \leq \sqrt{\varepsilon_\alpha} \rightarrow 0 \quad \text{and} \quad \|y_\alpha^* - x_\alpha^*\| \leq \sqrt{\varepsilon_\alpha} \rightarrow 0.$$

By $x_\alpha \xrightarrow{w^*} x^{***}$, we have $y_\alpha^* \xrightarrow{w^*} x^{***}$. ■

Remark. Clearly, $\partial f(x) \subset \partial\hat{f}(x)$ for all $x \in D$. This and Theorem 3.1 imply that for all $x \in D$,

$$\partial\hat{f}(x) = w^* - \lim_{y \rightarrow x} \partial f(y).$$

PROPOSITION 3.1. *Let f be a continuous convex function defined on an open convex subset D of X . Then $\partial\hat{f}(x) = w^* - \text{cl}_{X^{***}}(\partial f(x))$ if and only if for each weak neighborhood W of 0 in X^* there is $\delta > 0$ such $\partial f(B_X(x, \delta)) \subset \partial f(x) + W$.*

Proof. The “if” part, it is easy to verify $w^*\text{-cl}_{X^{***}}(\partial f(x)) \subset \partial \hat{f}(x)$. Suppose that there is $x_0^{***} \in \partial \hat{f}(x)$ such that $x_0^{***} \notin w^*\text{-cl}_{X^{***}}(\partial f(x))$. By the separation theorem, there are $x_0^{**} \in X^{**}$ and $r \in R$ such that

$$\langle x_0^{***}, x_0^{**} \rangle > r > \sup\{\langle x_0^{**}, x^* \rangle; x^* \in \partial f(x)\}.$$

Let $W = \{x^* \in X^*; \langle x_0^{**}, x^* \rangle < k\}$, where $k = r - \sup\{\langle x_0^{**}, x^* \rangle; x^* \in \partial f(x)\}$. Then W is a weak neighborhood of 0 in X^* . Hence there is $\delta > 0$ such that $B_X(x, \delta) \subset D$ and $\partial f(B_X(x, \delta)) \subset \partial f(x) + W$. By Theorem 3.1, there is a net $\{x_\alpha\}$ in D and a net $\{x_\alpha^*\}$ in X^* such that

$$x_\alpha^* \in \partial f(x_\alpha), \quad x_\alpha \rightarrow x \quad \text{and} \quad x_\alpha^* \xrightarrow{w^*} x_0^{***}.$$

Without loss of generality we can assume that $\{x_\alpha\} \subset B_X(x, \delta)$. Hence for each index α , $x_\alpha^* \in \partial f(x) + W$, and so $\langle x_0^{**}, x_\alpha^* \rangle < r$. This and $x_\alpha^* \xrightarrow{w^*} x_0^{***}$ imply $\langle x_0^{***}, x_0^{**} \rangle \leq r$, a contradiction. Hence $\partial \hat{f}(x) = w^*\text{-cl}_{X^{***}}(\partial f(x))$. For the “only if” part, it suffices to show that for any weak neighborhood W of 0 of the form

$$W = \{x^* \in X^*; |\langle x_i^{**}, x^* \rangle| < \varepsilon, x_i^{**} \in X^{**}, i = 1, \dots, n \text{ and } \varepsilon > 0\},$$

there is $\delta > 0$ such that $\partial f(B_X(x, \delta)) \subset \partial f(x) + W$. Since $\partial \hat{f}$ is $\|\cdot\|$ - w^* upper semicontinuous at $x \in D \subset \text{int}_{X^{**}}(\text{dom}(\hat{f}))$, there is $\delta > 0$ such that $\partial \hat{f}(B_{X^{**}}(x, \delta)) \subset \partial \hat{f}(x) + \frac{1}{2}\hat{W}$, where

$$\hat{W} = \{x^{***} \in X^{***}; |\langle x_i^{***}, x_i^{**} \rangle| < \varepsilon, i = 1, \dots, n\}.$$

Notice that $\partial \hat{f}(x) = w^*\text{-cl}_{X^{***}}(\partial f(x)) \subset \partial f(x) + \frac{1}{2}\hat{W}$, $\frac{1}{2}\hat{W} + \frac{1}{2}\hat{W} \subset \hat{W}$ and $\partial f(B_X(x, \delta)) \subset \partial \hat{f}(B_X(x, \delta))$. One has $\partial f(B_X(x, \delta)) \subset \partial f(x) + W$. ■

It is worth comparing Proposition 3.1 with Theorem 2.1 in [13].

COROLLARY 3.1. *Let f be a continuous convex function defined on an open convex subset D of X , and $x \in D$. Then $\partial \hat{f}(x) = \partial f(x)$ if and only if $\partial f(x)$ is a weak compact subset of X^* and ∂f is $\|\cdot\|$ - w upper semicontinuous at x .*

Proof. The “only if” part is trivial. It suffices to show that “if” part. By the weak compactness of $\partial f(x)$, we have $w^*\text{-cl}_{X^{***}}(\partial f(x)) = \partial f(x)$. This and Proposition 3.1 imply $\partial \hat{f}(x) = \partial f(x)$. ■

The following theorem is an immediate result of Corollary 3.1.

THEOREM 3.2. *Let f be a continuous convex function defined on an open convex subset D of X , and $x \in D$. Then f is very differentiable at x if and only if $\partial f(x)$ is a singleton and ∂f is $\|\cdot\|$ - w upper semicontinuous at x .*

Theorem 3.2 explains that, for a continuous convex function f , the very differentiability of f is between the Gateaux differentiability of f and the Frechet differentiability of f . Moreover, when X is a nonreflexive Banach space, in general, the very differentiability of f is not any kind of the β -differentiability of f .

LEMMA 3.1. *Let K be a weak* compact convex subset of X^* , and ∂p_K be $\|\cdot\|$ -w upper semicontinuous at $x_0 \in D$. Assume that a net $\{x_\alpha^*\}_{\alpha \in \Lambda}$ in K satisfies $\langle x_\alpha^*, x_0 \rangle \rightarrow p_K(x_0)$. Then there is a subnet $\{y_\beta^*\}_{\beta \in \Delta}$ of $\{x_\alpha^*\}$ and a net $\{z_\beta^*\}_{\beta \in \Delta}$ in $\partial p_K(x_0)$ such that $y_\beta^* - z_\beta^* \xrightarrow{w} 0$.*

Proof. Let $\varepsilon_\alpha = p_K(x_0) - \langle x_\alpha^*, x_0 \rangle$; then $\varepsilon_\alpha \geq 0$ and $\varepsilon_\alpha \rightarrow 0$. For each $h \in X$,

$$\begin{aligned} \langle x_\alpha^*, h \rangle &= \langle x_\alpha^*, x_0 + h \rangle - \langle x_\alpha^*, x_0 \rangle \leq p_K(x_0 + h) - \langle x_\alpha^*, x_0 \rangle \\ &= p_K(x_0 + h) - p_K(x_0) + \varepsilon_\alpha. \end{aligned}$$

Hence $x_\alpha^* \in \partial_{\varepsilon_\alpha} p_K(x_0)$. By the Brondsted-Rockafellar theorem, there is a net $\{x_\alpha\}_{\alpha \in \Lambda}$ in X and a net $\{a_\alpha^*\}_{\alpha \in \Lambda}$ in X^* such that

$$a_\alpha^* \in \partial p_K(x_\alpha), \|x_\alpha - x_0\| \leq \sqrt{\varepsilon_\alpha} \rightarrow 0 \text{ and } \|a_\alpha^* - x_\alpha^*\| \leq \sqrt{\varepsilon_\alpha} \rightarrow 0. \quad (3)$$

Let $\Delta = \{(\alpha, U); \alpha \in \Lambda \text{ and } U \text{ is a weak neighborhood of } 0 \text{ in } X^*\}$, and define a partial order on Δ as follows: for $(\alpha_1, U_1), (\alpha_2, U_2) \in \Delta$, $(\alpha_1, U_1) \leq (\alpha_2, U_2)$ if and only if $\alpha_1 \leq \alpha_2$ and $U_2 \subset U_1$. It is clear that (Δ, \leq) is a directed set. Since ∂p_K is $\|\cdot\|$ -w upper semicontinuous at x_0 , by (3), for each $(\alpha, U) \in \Delta$, there is $\varphi(\alpha, U) \in \Lambda$ such that $\alpha \leq \varphi(\alpha, U)$ and $a_{\varphi(\alpha, U)}^* \in \partial p_K(x_0) + U$. Hence there is $b_{\varphi(\alpha, U)}^* \in \partial p_K(x_0)$ such that $a_{\varphi(\alpha, U)}^* - b_{\varphi(\alpha, U)}^* \in U$. Denote $x_{\varphi(\alpha, U)}^*$ and $b_{\varphi(\alpha, U)}^*$ by $y_{(\alpha, U)}^*$ and $z_{(\alpha, U)}^*$, respectively. It is easy to verify that $\{y_{(\alpha, U)}^*\}_{(\alpha, U) \in \Delta}$ is a subnet of $\{x_\alpha^*\}_{\alpha \in \Lambda}$, $\{z_{(\alpha, U)}^*\}_{(\alpha, U) \in \Delta}$ is a net in $\partial p_K(x_0)$, and $y_{(\alpha, U)}^* - z_{(\alpha, U)}^* \xrightarrow{w} 0$. ■

PROPOSITION 3.2. *Let K be a weak* compact convex subset of X^* , and $x_0 \in X$. The following properties are equivalent:*

(i) *Each point x^* in $\partial p_K(x_0)$ is a weak*-weak continuous point of K (i.e., each net in K is weakly convergent to x^* whenever it is weak* convergent to x^*).*

(ii) *Each net $\{x_\alpha^*\}$ in K with $\langle x_\alpha^*, x_0 \rangle \rightarrow p_K(x_0)$ has a subnet which is weakly convergent to some point in $\partial p_K(x_0)$.*

(iii) $\partial \hat{p}_K(x_0) = \partial p_K(x_0)$.

Proof. (i) \Rightarrow (ii). By weak* compactness of K , each net $\{x_\alpha^*\}$ in K with $\langle x_\alpha^*, x_0 \rangle \rightarrow p_K(x_0)$ has a subnet which is weak* convergent to some x^* in K . It follows that $\langle x^*, x_0 \rangle = p_K(x_0)$. By [10, Lemma 5.10], $x^* \in \partial p_K(x_0)$. This and (i) imply that $\{x_\alpha^*\}$ has a subnet which is weakly convergent to x^* .

(ii) \Rightarrow (iii). It is clear from (ii) that $\partial p_K(x_0)$ is weakly compact. Suppose that $\partial \hat{p}_K(x_0) \neq \partial p_K(x_0)$. By Corollary 3.1, there is a weak open set W including $\partial p_K(x_0)$, a sequence $\{x_n\}$ in X and a sequence $\{x_n^*\}$ in X^* such that $x_n \rightarrow x_0$, $x_n^* \in \partial p_K(x_n)$ and $x_n^* \notin W$ for all n . Since

$$\begin{aligned}\langle x_n^*, x_0 \rangle &= \langle x_n^*, x_n \rangle + \langle x_n^*, x_0 - x_n \rangle \\ &= p_K(x_n) + \langle x_n^*, x_0 - x_n \rangle \rightarrow p_K(x_0),\end{aligned}$$

without loss of generality we can assume that there is $x^* \in K$ such that $x_n^* \xrightarrow{w} x^*$. It follows that $x^* \notin W$ and $\langle x^*, x_0 \rangle = p_K(x_0)$. This and Lemma 5.10 in [10] implies $x^* \in \partial p_K(x_0)$. This contradicts $\partial p_K(x_0) \subset W$.

(iii) \Rightarrow (i). Let x_0^* be any point in $\partial p_K(x_0)$ and $\{x_\beta^*\}$ be any net in K such that $x_\beta^* \xrightarrow{w^*} x_0^*$. For any subnet $\{y_\alpha^*\}$ of $\{x_\beta^*\}$, by Lemma 3.1 and the weak compactness of $\partial p_K(x_0)$, $\{y_\alpha^*\}$ has a subnet which is weakly convergent. This implies $x_\beta^* \xrightarrow{w^*} x_0^*$. ■

Let K be a weak* compact subset of X^* and K_0 be the weak* closed convex hull of K . Notice that for each $x \in X$ there is $x^* \in K$ such that

$$\langle x^*, x \rangle = \sup\{\langle y^*, x \rangle; y^* \in K\} = \sup\{\langle y^*, x \rangle; y^* \in K_0\}.$$

One has the following corollaries.

COROLLARY 3.2. *Let K be a weakly compact subset of X^* . Then $\partial \hat{p}_K(x) = \partial p_K(x)$ for all $x \in X$.*

COROLLARY 3.3. *Let K be a weak* compact convex subset of X^* and $x_0 \in X$. Then p_K is very differentiable at x_0 if and only if each net $\{x_\alpha^*\}$ in K with $\langle x_\alpha^*, x_0 \rangle \rightarrow p_K(x_0)$ is weakly convergent.*

COROLLARY 3.4. *Let K be a weak* compact subset of X^* and $x_0 \in X$. If $\partial \hat{p}_K(x_0) = \partial p_K(x_0)$, there are $x_1 \in X$ and $x_0^* \in K$ such that each net $\{x_\alpha^*\}$ in K with $\langle x_\alpha^*, x_i \rangle \rightarrow \langle x_0^*, x_i \rangle$ ($i = 0, 1$) is weakly convergent to x_0^* . In particular, x_0^* is a weak*-weak continuous point of K .*

Proof. Let K_0 be the weak* closed convex hull of K . By Corollary 3.1, $\partial p_K(x_0)$ is a weakly compact convex subset of K_0 . Hence there are $x_1 \in X$ and $x_0^* \in \partial p_K(x_0)$ such that x_1 strongly exposes $\partial p_K(x_0)$ at x_0^* . Notice that $\partial p_K(x_0)$ is an extremal subset of K_0 and x_0^* is an extremal point of $\partial p_K(x_0)$. x_0^* is an extremal point of K_0 . Hence $x_0^* \in K$ (see [2, Theorem 36.10]). Let $\{x_\alpha^*\}$ be any net in K with $\langle x_\alpha^*, x_i \rangle \rightarrow \langle x_0^*, x_i \rangle$ ($i = 0, 1$). By Proposition 3.2, it is sufficient to prove that each subnet of $\{x_\alpha^*\}$ is weakly convergent to x_0^* whenever the subnet is weakly convergent. Assume that a subnet $\{y_\beta^*\}$ of $\{x_\alpha^*\}$ is weakly convergent. By Lemma 3.1, there is a subnet $\{a_\gamma^*\}$ of $\{y_\beta^*\}$ and a net $\{b_\gamma^*\}$ in $\partial p_K(x_0)$ such that $a_\gamma^* - b_\gamma^* \xrightarrow{w} 0$. This

implies $\langle b_\gamma^*, x_1 \rangle \rightarrow \langle x_0^*, x_1 \rangle$. Hence $\|b_\gamma^* - x_0^*\| \rightarrow 0$. It follows that $\{y_\beta^*\}$ is weakly convergent to x_0^* . ■

4. A CHARACTERIZATION OF ASPLUND SPACES

In this section we will give an characterization of Asplund spaces which is similar to Theorem 2.7 in [4]. We need the following lemma.

LEMMA 4.1 ([4, Theorem 1.3]). *Let Φ be a minimal w^* -cusco from a complete metric space A into the bidual space X^{**} of a Banach space X . Assume that the set*

$$E = \{x \in A; \Phi(x) \subset w^*\text{-cl}_{X^{**}}(\Phi(x) \cap X)\}$$

*is a residual subset of A (i.e., E includes a dense G_δ subset of A). Then Φ is single-valued and upper semicontinuous with respect to the norm topology of X^{**} at each point in a dense G_δ subset of A .*

PROPOSITION 4.1. *X is an Asplund space if and only if for each bounded subset K of X^* there is a metric ρ on K such that every bounded subset of K admits weak* slices of arbitrarily small ρ -diameter and the topology induced by ρ on K is stronger than or equal to the weak topology.*

Proof. The “only if” part is immediate from Theorem 2.32 in [10]. To prove the “if” part, it suffices to show that, for each open convex subset D of X and each continuous convex function f defined on D , f is Frechet differentiable on a dense subset of D (because the set of all points of D where f are Frechet differentiable is a G_δ subset of D). Since the subdifferential mapping ∂f is locally bounded, for each $x \in D$ and each neighborhood U of x there is an open convex neighborhood V of x such that $V \subset U \cap D$ and $\partial f(V)$ is bounded. Hence there is a metric ρ on $\partial f(V)$ such that every bounded subset of $\partial f(V)$ admits weak* slices of arbitrarily small ρ -diameter and the topology induced by ρ on $\partial f(V)$ is stronger than or equal to the weak topology. For each natural number n , let $G_n = \{x \in V; \text{there is an open subset } W \text{ of } V \text{ such that } x \in W \text{ and } \rho - \text{diam}(\partial f(W)) < \frac{1}{n}\}$. Similar to the proof of [10, Theorem 2.30], we have that $G = \bigcap_{n=1}^\infty G_n$ is a dense G_δ subset of V . It is clear that for each $x \in G$, $\partial f(x)$ is a singleton and ∂f is $\|\cdot\|_\rho$ upper semicontinuous at x , and so ∂f is $\|\cdot\|_w$ upper semicontinuous at x . It is sufficient to show that f is Frechet differentiable on a dense subset of V . Let T be the restriction of ∂f to G . Then T is a $\|\cdot\|_w$ continuous single-valued mapping from G into X^* . Hence T is a minimal w -cusco from G into X^* . Since a dense G_δ subset of a complete metric space is itself completely metrizable ([5, p.

96]), by Lemma 4.1 there is a dense G_δ subset A of G such that for each $x \in A$, T is $\|\cdot\| \cdot \|\cdot\|$ upper semicontinuous at x . This and Lemma 2.5 in [4] imply that ∂f , as a set-valued mapping from V into X^* , is single-valued and $\|\cdot\| \cdot \|\cdot\|$ upper semicontinuous at each point of A . Hence f is Frechet differentiable at each point of the dense subset A of V . ■

In view of Proposition 4.1 the following problem is natural.

Is X an Asplund space if every bounded subset of X^* admits arbitrarily small weak* slices with respect to the weak topology (i.e., for any weak neighborhood W of 0 in X^* and any bounded subset K of X^* there are $x \in X$ and $\alpha > 0$ such that $S(x, K, \alpha) - S(x, K, \alpha) \subset W$, where $S(x, K, \alpha) = \{x^* \in K; \langle x^*, x \rangle > \sup\{\langle y^*, x \rangle; y^* \in K\} - \alpha\}$)?

Consider the James Tree space JT which is a separable Banach space with nonseparable dual space. In addition every infinite dimensional subspace of JT contains a copy of l_2 (see [3]). Hence JT is not an Asplund space and contains no copy of l_1 . Notice that X contains no copy of l_1 if and only if every bounded subset K of X^* admits arbitrarily small weak* slices with respect to the weak topology (see [14, pp. 351–352]). It follows that JT is not an Asplund space and every bounded subset of JT^* admits arbitrarily small weak* slices with respect to the weak topology.

G. Godefroy [7] really gave the following characterization of an Asplund space:

X is an Asplund space if and only if each equivalent norm on X has a very differentiable point in X (see the proof of Theorem II.2 and Remark II.3 in [7]).

It follows directly from Godefroy's result and Corollary 3.3 that X is an Asplund space if and only if for each weak* compact convex subset K of X^* there are $x \in X$ and $x^* \in K$ such that each net $\{x_\alpha^*\}$ is weakly convergent to x^* whenever $\langle x_\alpha^*, x \rangle \rightarrow \langle x^*, x \rangle = p_K(x)$. But the existence of weak*-weak continuous points of each weak* compact convex subset of X^* does not imply that X is an Asplund space. Indeed, Ghoussoub, Maurey, and Schachermayer [6] have shown that the existence of weak*-norm continuous points of each weak* compact convex subset of X^* does not imply that X is an Asplund space.

The following result is an equivalent form of Theorem 3.6 in Moors [8].

X is an Asplund space if and only if, for each weak* compact subset K of X^* , the set of weak*-weak continuous points of K is residual in (K, w^*) .

In view of above-presented results and Corollary 3.4, we present the following problem:

Is X an Asplund space if, for each equivalent norm $\|\cdot\|$ (each continuous seminorm p) on X , there is $x \in X$ such that $\partial \|\cdot\|^\wedge(x) = \partial \|\cdot\|(x)(\partial \hat{p}(x) = \partial p(x))$?

Note added in Proof. Giles [15] answered the above problem “yes” sometime after this paper was submitted.

ACKNOWLEDGMENTS

The author expresses his gratitude to the referee for many helpful suggestions which improve this paper. This work was supported by the ABSF of Yunnan Province, P. R. China.

REFERENCES

1. E. Asplund, Frechet differentiability of convex function, *Acta Math.* **88** (1968), 35–46.
2. S. K. Berberian, “Lectures in Functional Analysis and Operator Theory,” Springer-Verlag, New York, 1974.
3. J. Diestel and J. J. Uhl, “Vector Measures,” Math. Surveys 15, Amer. Math. Soc., Providence, 1977.
4. J. R. Giles, P. S. Kenderov, W. B. Moors, and S. D. Sciffer, Generic differentiability of convex functions on the dual of a Banach space, *Pacific J. Math.* **172** (1996), 413–433.
5. J. L. Kelley and I. Namioka, “Linear Topological Spaces,” Springer, New York, 1976.
6. N. Ghoussoub, B. Maurey, and W. Schachermayer, Geometrical implications of certain infinite dimensional decompositions, *Trans. Amer. Math. Soc.* **317** (1990), 541–584.
7. G. Godefroy, Metric characterizations of first Baire class linear forms and octahedral norms, *Studia Math.* **95** (1989), 1–15.
8. W. B. Moors, The relationship between Goldstine’s theorem and the convex point of continuity property, *J. Math. Anal. Appl.* **188** (1994), 819–832.
9. I. Namioka and R. R. Phelps, Banach spaces which are Asplund spaces, *Duke Math. J.* **42** (1975), 735–750.
10. R. R. Phelps, Convex functions, in “Monotone Operators and Differentiability,” Lecture Notes in Mathematics, No. 1364, Springer-Verlag, New York, 1989.
11. C. Stegall, The duality between Asplund spaces and spaces with the Radon-Nikodym property, *Israel J. Math.* **29** (1978), 408–412.
12. N. K. Thakare, Geometry of Banach spaces: Rough and very smooth norms, Asplund spaces and related topics, in “Modern Analysis and Its Applications,” Prentice-Hall of India, New Delhi, 1986.
13. W. B. Moors, Generic continuity of restricted upper semicontinuous set-valued mappings, *Set-Valued Anal.* **4** (1996), 25–39.
14. X. T. Yu, “Selected Topics on Banach Spaces,” East China Normal Univ. Publishing House, Shanghai, 1992.
15. J. R. Giles, Comparable characterizations of two classes of Banach spaces by subdifferentials, *Bull. Austral. Math. Soc.* **58** (1998), 155–158.